



AUTHOR: PLEASE COMPLETE

Read & Corrected by

Date

RESEARCH BRIEF

Crithidia fasciculata: A Test For Genetic Exchange

Jan Votýpka,*† Dan S. Ray,‡ and Julius Lukeš*§¹

*Institute of Parasitology, Czech Academy of Sciences, Branišovská 31, 37005 České Budějovice, Czech Republic;

†Department of Parasitology, Faculty of Sciences, Charles University, Prague, Czech Republic;

‡Molecular Biology Institute and Department of Microbiology and Molecular Genetics, University of California, Los Angeles, California, U.S.A.; and §Faculty of Biology, University of South Bohemia, České Budějovice, Czech Republic

Votýpka, J., Ray D. S., and Lukeš J. 2001. *Crithidia fasciculata*: A test for genetic exchange. *Experimental Parasitology* 00, 000–000.

© 2001 Academic Press

Index Descriptors and Abbreviations: *Crithidia*; Kinetoplastida; sex; drug resistance; transformation; mosquito; transmission; recombination.

Genetic exchange between cells of *Trypanosoma brucei* that occurs in the tsetse fly vector (Jenni *et al.* 1986) is an important part of the life cycle that contributes significantly to genetic diversity of these human pathogens (Gibson and Stevens 1999). Indirect observations have also indicated the presence of sexuality during the development of *Trypanosoma cruzi* (Bogliolo *et al.* 1996; Stothard *et al.* 1999) and *Trypanosoma carassii* (Zajfček 1991). However, a thorough search for this process in related *Leishmania* failed to detect the recombination process in its vector (Panton *et al.* 1991), although rare natural hybrids have been reported (Belli *et al.* 1994; Banuls *et al.* 1997). A better understanding of genetic exchange that apparently has important implications for pathogenicity and epidemiology of sleeping sickness and Chagas disease would also provide us with a novel way of combining the limited number of markers (Clayton 1999) currently available for genetic manipulations of Kinetoplastida. *Crithidia fasciculata* is a monoxenic model trypanosomatid, the Steinert and UC strains of which have existed in culture for decades. For *C. fasciculata*, a relatively high frequency of appearance of mutants with multiple resistance has

been reported (Hughes *et al.* 1982). It was also observed that a mixed growth of its mutants resistant to different drugs resulted in the selection of double-resistant clones, suggesting the existence of some type of genetic exchange (Glassberg *et al.* 1985). Moreover, close apposition of the *C. fasciculata* cells during the process of flagellar adherence was reminiscent of cell fusion of other types of recombination (Hughes *et al.* 1983). To our knowledge, these results were never verified and no study is available that would search for sexuality of *Crithidia* in its mosquito host. In this work, using a highly sensitive approach, we have attempted to detect genetic exchange in this flagellate both in the insect vector and in culture.

We have used the UC strain of *C. fasciculata* genetically modified by the disruption of one allele of the single-copy ribonuclease H1 gene (*RNHI*) via targeted insertion of the neomycin (*NEO* strain) or G418 (*HYG* strain) drug-resistance cassettes as described (Ray and Hines 1995). The *RNHI* gene encodes both nuclear and mitochondrial isoforms of ribonuclease H (Engel *et al.* 2001) and seems to be nonessential, since the double knock-out strain was viable (Ray and Hines 1995). With the *C. fasciculata* strains carrying either the *NEO* or the *HYG* markers in one of their *RNHI* alleles, we had a powerful tool in our hands to search for a possible genetic cross bearing both markers in its chromosomes. Since genetic recombination in *T. brucei* has been convincingly demonstrated to occur in the vector, we have attempted to infect different mosquito species with our strains. With their decades-long cultivation in the culture form only, it was of interest to determine whether after innumerable generations the parasite did not lose its potential to develop in the insect host. Among mosquito species listed by Wallace (1966) as natural hosts of *C. fasciculata*, we were able to obtain the cultures of *Culex pipiens quinquefasciatus* and *Aedes aegypti*, and we have included *Culex pipiens molestus* in our infection experiments.

The *NEO* and *HYG* strains tolerated concentrations of G418 and

¹To whom correspondence should be addressed. Fax: +420-38-5300388. E-mail: jula@paru.cas.cz.



hygromycin up to 0.25 and 0.5 mg/ml, respectively. The *NEO* strain was highly sensitive to hygromycin (lethal concentration 20 µg/ml), whereas the *HYG* strain died after the addition of 10 µg/ml of G418. Both strains were cultivated in BHI medium (Difco) containing hemin (20 µg/ml), streptomycin (100 µg/ml), penicillin (500 U/ml), and ampicillin (100 µg/ml) at 24°C. The *C. fasciculata* UC wild-type (*WT*) strain was used in control experiments. The IVth larval instars of *C. p. quinquefasciatus* (strain "H;" originating from Hyderabad, India; colonized in 1980), *C. p. molestus* (strain "M;" originating from Mosul, Iraq; colonized in 1989), *C. p. molestus* (strain "CB;" originating from České Budějovice, Czech Republic; colonized in 1986), and *A. aegypti* (an old laboratory strain without particular identification) obtained from established colonies were transferred into water containing 100 µg/ml of ampicillin to minimize bacterial contamination and were reared to adulthood. After emergence, 7- to 10-day-old adults were offered a sterile cotton pad soaked with BHI medium supplemented with 10% sucrose solution containing the *C. fasciculata* *NEO* and/or *HYG* strains (Table I) in which the flagellates remained alive for more than 10 h. For the following 4 to 11 days (depending on the experiment) the mosquitoes were fed with a sterile 10% sucrose solution provided in a cotton pad. After aspiration and subsequent immobilization by low temperature, the adults were immersed in 70% ethanol and washed in sterile phosphate-buffered saline (PBS), and under the dissection microscope their hindgut and rectal ampula were carefully removed. After being checked for the flagellates under light microscope, the tissue was homogenized by gentle squeezing with a cover slip, and the homogenate was then dissolved in 100 µl of sterile PBS. Two milliliters of BHI medium containing either 50 µg/ml G418 or 100 µg/ml hygromycin each were inoculated with 10 µl of the above homogenate, and 30 µl of it was added to BHI medium containing both agents. The same volumes were applied on plates with solid BHI media containing 1% agar supplemented with 70 µg/ml G418 and/or 150 µg/ml hygromycin (Table I). The inoculated media were kept for up to 10 days in an orbital shaker at 24°C and checked daily for flagellates. The agar plates were sealed with parafilm and placed into a thermostat at 28°C for up to 1 month. They were regularly checked for pinhead-sized colonies originating from a single cell.

Prior to exposure to *C. fasciculata*, all mosquito cultures used were shown both by microscopy and by cultivation to be negative for any

flagellates. Next, the *C. fasciculata* strains (*NEO*, *HYG*, and *WT*) separately fed to 10-day-old adults of both sexes of the tested mosquito species were proved to be equally infective. The level of infection was different in individual mosquitoes but we have not investigated possible correlation between the flagellate and the vector species and/or sex. Heavy infections, with the intestinal contents of the host teeming with flagellates adhered to the intestinal wall, occurred in each case. Uninfected mosquitoes were then offered a pad soaked with the cultures of both the *NEO* and the *HYG* strains. After dissection, the intestinal contents of each mosquito were used for the inoculation of BHI media and agar plates containing appropriate amounts of G418 (column *NEO*), hygromycin (column *HYG*), and both G418 and hygromycin (column *NEO/HYG*) (Table I). The problem of bacterial contamination was largely solved by rearing of the insects in the antibiotics-containing water.

As summarized in Table I, all inoculations in liquid media with a single inhibitory agent resulted in the appearance of flagellates within 2 to 5 days. The same was true for the agar plates on which, however, colonies were spotted after 2- to 3-week-long incubation. Generally, lower numbers of colonies appeared on the G418 plates than on the hygromycin plates in the same experiment. We explain this discrepancy by the lower number of G418-resistant flagellates streaked on the plates, since these cells grow somewhat slower than the hygromycin-resistant cells. The high total number of colonies that grew on plates with a single selective agent can in fact be tripled, since only 10 µl of the intestinal homogenate was used for their inoculation compared to 30 µl applied on the plates and liquid media containing both drugs. Still, under these experimental conditions we have never observed cells with both the *NEO* and the *HYG* cassettes in their genome enabling them to grow in media containing both selection agents.

An experiment with the same design that used flagellates stably transformed with genes conferring drug resistance provided further proof for genetic exchange in *T. brucei* (Gibson and Whittington 1993). Due to the tagging of a single-copy gene, our experiments were even more sensitive than those of Gibson and Whittington, who tagged only one allele of a tubulin gene present on two or more chromosomes, and in principle we should thus be able to detect every Mendelian cross.

We have also performed a mixed growth of the *NEO* and *HYG* strains in culture. Depending on the experiment, the strains with different drug

TABLE I
Results of Mixed Growth of Two *C. fasciculata* Strains (*NEO* and *HYG*) in Natural Hosts

Species	Mosquitoes		Liquid medium			Solid medium		
	Sex	Number	<i>NEO</i>	<i>HYG</i>	<i>NEO/HYG</i>	<i>NEO</i>	<i>HYG</i>	<i>NEO/HYG</i>
<i>Aedes aegypti</i>	M	5	+	+	-	18	72	0
	F	24	+	+	-	121	520	0
<i>Culex p. quinquefasciatus</i>	M	4	+	+	-	38	72	0
	F	4	+	+	-	51	124	0
<i>Culex p. molestus</i> "M"	M	1	+	+	-	12	28	0
	F	11	+	+	-	227	520	0
<i>Culex p. molestus</i> "CB"	M	5	+	+	-	159	162	0
	F	11	+	+	-	763	731	0
Total		65	+	+	-	1389	2229	0

Note. The (+) means intensive growth in the liquid medium after 5 days of incubation. Numbers in the *NEO* and *HYG* columns (solid medium) indicate number of colonies on plates after 21 days of incubation.

TABLE II

Results of Mixed Growth of Two *C. fasciculata* Strains (*NEO* and *HYG*) in Liquid Medium

Exp. No.	Days of growth	Cell density per ml	Liquid medium			Solid medium		
			<i>NEO</i>	<i>HYG</i>	<i>NEO/HYG</i>	<i>NEO</i>	<i>HYG</i>	<i>NEO/HYG</i>
1	3	3.1×10^6	+	+	—	160	430	0
2	4	2.5×10^7	+	+	—	1100	2100	0
3	4	3.0×10^7	+	+	—	120	330	0
4	5	1.8×10^8	+	+	—	150	380	0
5	5	2.1×10^8	+	+	—	60	150	0
Total			+	+	—	1590	3390	0

Note. The (+) means intensive growth in the liquid medium after 5 days of incubation. Numbers in the *NEO* and *HYG* columns (solid medium) indicate number of colonies on plates after 21 days of incubation.

resistance were cocultivated for 3 to 5 days and then transferred to the liquid media or plates containing both drugs (Table II). Although a high number of colonies grew on the G418- or hygromycin-containing plates, the media with both selective agents remained negative (Table II).

We did not find any support for the existence of genetic exchange in *C. fasciculata*, both in culture and in the insect host. The occurrence of such a process in cultured promastigotes of this species suggested by Glassberg *et al.* (1985) has to be considered unlikely since it was neither detected by us nor described in any flagellate studied so far (Gibson and Stevens 1999). Negative results were reported also for *Leishmania*, where the authors speculated that the sensitivity of their test may not have been sufficient to detect very rare sexual events (Panton *et al.* 1991). The stable chromosomal markers used in our experiments represent an even more sensitive tool but they failed to prove the existence of such a process in *Crithidia*. Although more data are needed, in the light of our data we propose that genetic exchange is a process absent or extremely rare in the life cycle of *C. fasciculata*.

ACKNOWLEDGMENTS

(We thank Dr. Jiří Olejníček (Institute of Parasitology, České Budějovice) for providing the IVth instar mosquito larvae. This work was supported by grant 204/00/1212 from the Grant Agency of the Czech Republic to J.L., grant Z6022909 from the Czech government, and grant AI 45536 from the National Institutes of Health to D.S.R.)

REFERENCES

Banuls, A. L., Guerrini, F., LePont, F., Barrera, C., Espinel, I., Guderian, R., Echeverria, R., and Tibayrenc, M. 1997. Evidence for hybridization by multilocus enzyme electrophoresis and random amplified polymorphic DNA between *Leishmania braziliensis* and *Leishmania*

panamensis/guyanensis in Ecuador. *Journal of Eukaryotic Microbiology* **44**, 408–411.

Belli, A. A., Miles, M. A., and Kelly, J. M. 1994. A putative *Leishmania panamensis/Leishmania braziliensis* hybrid is a causative agent of human cutaneous leishmaniasis in Nicaragua. *Parasitology* **109**, 435–442.

Bogliolo, A. R., Lauria-Pires, L., and Gibson, W. 1996. Polymorphism in *Trypanosoma cruzi*: Evidence of genetic recombination. *Acta Tropica* **61**, 31–40.

Clayton, C. E. 1999. Genetic manipulation of Kinetoplastida. *Parasitology Today* **15**, 372–378.

Engel, M. L., Hines, J. C., and Ray, D. S. 2001. The *Crithidia fasciculata* RNH1 gene encodes both nuclear and mitochondrial isoforms of RNase H. *Nucleic Acids Research* **29**, 725–731.

Gibson, W., and Whittington, H. 1993. Genetic exchange in *Trypanosoma brucei*: Selection of hybrid trypanosomes by introduction of genes conferring drug resistance. *Molecular and Biochemical Parasitology* **60**, 19–26.

Gibson, W., and Stevens, J. 1999. Genetic exchange in the Trypanosomatidae. *Advances in Parasitology* **43**, 1–63.

Glassberg, J., Miyazaki, L., and Rifkin, M. R. 1985. Isolation and partial characterization of mutants of the trypanosomatid *Crithidia fasciculata* and their use in detecting genetic recombination. *Journal of Protozoology* **32**, 118–125.

Hughes, D. E., Schneider, C. A., and Simpson, I. 1982. Isolation and characterization of drug resistant mutants of *Crithidia fasciculata*. *Journal of Parasitology* **68**, 642–649.

Hughes, D. E., Simpson, L., and Schneider, C. A. 1983. Flagellar adherence of *Crithidia fasciculata* cells in culture. *Journal of Protozoology* **30**, 635–641.

Jenni, L., Marti, S., Schweitzer, J., Betschart, B., LePage, R. W. F., Wells, J. M., Tait, A., Paindavoine, P., Pays, E., and Steinert, M. 1986. Hybrid formation between African trypanosomes during cyclical transmission. *Nature* **322**, 173–175.

Panton, L. J., Tesh, R. B., Nadeau, K. C., and Beverley, S. H. 1991. A test for genetic exchange in mixed infections of *Leishmania major*

- in the sand fly *Phlebotomus papatasi*. *Journal of Protozoology* **38**, 224–228.
- Ray, D. S., and Hines, J. C. 1995. Disruption of the *Crithidia fasciculata* RNH1 gene results in the loss of two active forms of the ribonuclease H. *Nucleic Acids Research* **23**, 2526–2530.
- Stothard, J. R., Frame, I. A., and Miles, M. A. 1999. Genetic diversity and genetic exchange in *Trypanosoma cruzi*: Dual drug-resistant “progeny” from episomal transformants. *Memorias do Instituto Oswaldo Cruz* **94**, 189–193.
- Wallace, F. G. 1966. The trypanosomatid parasites of insects and arachnids. *Experimental Parasitology* **18**, 124–193.
- Zajíček, P. 1991. *Trypanosoma* cf. *carassii*: The combination of malic enzyme patterns supports the theory of diploidy in trypanosomes. *International Journal for Parasitology* **21**, 753–755.

Received 2 November 2000; accepted with revision 28 August 2001